

A Comment on “Dynamics of Weak First Order Phase Transitions”

Geoffrey Harris
Enrico Fermi Institute
University of Chicago
Chicago, IL 60637, USA

Gerard Jungman
Physics Department
Syracuse University
Syracuse, NY 13244, USA
 (July 1995)

Nucleation and spinodal decomposition, as mechanisms for first-order transitions, have been studied for many years. A pertinent issue is whether or not there is a sharp distinction between these two processes. In Ref. [1], M. Gleiser analyzes a lattice model with one free parameter α which is tuned to control the strength of a first-order transition. He argues that there exists a critical value α_c distinguishing between strong and weak first-order transitions and suggests that only for $\alpha > \alpha_c$ does phase transformation proceed by nucleation. We believe that the numerical results in this work are correct, but will argue against the interpretation of the theory at α_c as a boundary between weak and strong phase transitions.

The phenomenology of phase transitions can be described by the Landau–Ginzburg Hamiltonian

$$\frac{H}{\theta} = \frac{1}{\theta} \left(\frac{1}{2} (\nabla \phi)^2 \pm m^2 \phi^2 + \frac{\lambda}{4} \phi^4 + \beta(t) \phi + \dots \right). \quad (1)$$

As the system is cooled, $\beta(t)$ changes sign, the minimum of the free-energy shifts discontinuously, and phase transformation occurs. In Ref. [1], the above Hamiltonian, with $\beta(t) \equiv 0$ and a positive coefficient of ϕ^2 , is adopted as a *lattice* Hamiltonian. This free energy has two degenerate minima; ϕ is placed throughout the lattice in the left-most minima and its relaxation (with second order Langevin dynamics) is studied; α controls the height of the hump between the two minima. At a certain non-zero value of α , the relaxation dynamics becomes critical. The size of the potential hump is interpreted as a measure of the strength of a first-order transition that would occur under cooling, implemented by a time-dependent parity-odd perturbation of the Hamiltonian. For $\alpha \gg \alpha_c$, the bubble free energy is large and nucleation should adequately describe the relaxation. It is then inferred that a different relaxation process may operate when $\alpha < \alpha_c$.

This description is misleading because it depends on the shape of the lattice (bare) potential. The physics of the phase transformation is modeled by the *coarse-grained* potential, related to the bare potential by renormalization. Criticality of the relaxation process depends only on the long-time equilibrium behavior. Thus, we only need the renormalization corrections for a classical statistical mechanical model at finite temperature θ , or equivalently, a quantum field theory with $\hbar = 1/\theta$. Explicitly, consider the one-loop correction to the quadratic term, which depends on the inverse lattice spacing Λ as

$$H_{ct} = \frac{1}{4\pi} 3\lambda\theta \ln \left(\frac{\Lambda^2}{\mu^2(m, \lambda\theta)} \right) \phi^2; \quad (2)$$

the finite part of the counterterm, μ , is determined by imposing renormalization conditions. The leading order Λ dependent piece reduces the barrier in the coarse-grained free energy by an additive correction. We anticipate that at some value of α , presumably α_c , the quadratic term and bump in the continuum effective potential vanish. This corresponds to the second-order phase transition in the ϕ^4 system, which is in the Ising universality class.

Indeed, with this correspondence in mind, simulations of the lattice Hamiltonian of Ref. [1] have been previously performed. In Ref. [2], Monte Carlo simulations are used to extract Ising critical exponents from this model. In Ref. [3], Monte Carlo and Langevin simulations are used to obtain the critical line of the lattice ϕ^4 model. The coupling χ of Ref. [3] is exactly the quantity $\lambda\theta_c$ of Ref. [1]; the coupling θ^{TC} of Ref. [3], which we distinguish by a superscript *TC*, equals $(\theta_c^2 - 1)/2$ in Ref. [1], with $\theta_c = (1 - 2\alpha^2/9\lambda)^{-1/2}$. In Ref. [1], criticality occurs at $(\lambda, \alpha) = (.1, .36)$. This corresponds to $(\chi, \theta^{TC}) = (.119, .202)$, which lies within statistical error on the critical line presented in Table II and Figure 8 of Ref. [3]. We have repeated these numerical simulations, using Langevin dynamics as in Ref. [3]. We find results consistent with those above. Thus the critical behavior in Ref. [1] is in the Ising universality class,

with vanishing coarse-grained barrier energy. It is not appropriately described in the continuum limit as a boundary between large and small barrier energies.

G.H. and G.J. were supported by DOE grant DE-FG02-85ER-40231 at Syracuse University. Additionally, G.H. was supported by DOE grant DE-FG02-90ER-40560 at the University of Chicago.

- [1] M. Gleiser, Phys. Rev. Lett. **73**, 3495 (1994).
- [2] A. Milchev, D.W. Heermann and K. Binder, J. Stat. Phys. **44**, 749 (1986).
- [3] R. Toral and A. Chakrabarti, Phys. Rev. **B42**, 2445 (1990).